

A non-genomic signaling pathway shut down by mating changes the estradiol-induced gene expression profile in the rat oviduct

Alexis Parada-Bustamante^{1,2}, Pedro A Orihuela^{3,4}, Mariana Ríos^{1,2}, Catherina A Cuevas³, Maria Lorena Oróstica³, Luis A Velásquez^{3,4}, Manuel J Villalón¹ and Horacio B Croxatto^{2,3,4}

¹Unidad de Reproducción y Desarrollo, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, 8331010 Santiago, Chile, ²Millennium Institute for Fundamental and Applied Biology, 7780272 Santiago, Chile, ³Laboratorio de Inmunología de la Reproducción, Facultad de Química y Biología, Universidad de Santiago de Chile, 9170022 Santiago, Chile and ⁴Centro para el Desarrollo de la Nanociencia y la Nanotecnología (CEDENNA), 9170124 Santiago, Chile

Correspondence should be addressed to A Parada-Bustamante who is now at School of Medicine, Institute of Maternal and Child Research, University of Chile, Santa Rosa 1234, Segundo Piso, Casilla 226-3, Santiago, Chile; Email: aparadab@med.uchile.cl

Abstract

Estradiol (E₂) accelerates oviductal egg transport through intraoviductal non-genomic pathways in unmated rats and through genomic pathways in mated rats. This shift in pathways has been designated as intracellular path shifting (IPS), and represents a novel and hitherto unrecognized effect of mating on the female reproductive tract. We had reported previously that IPS involves shutting down the E₂ non-genomic pathway up- and downstream of 2-methoxyestradiol. Here, we evaluated whether IPS involves changes in the genomic pathway too. Using microarray analysis, we found that a common group of genes changed its expression in response to E₂ in unmated and mated rats, indicating that an E₂ genomic signaling pathway is present before and after mating; however, a group of genes decreased its expression only in mated rats and another group of genes increased its expression only in unmated rats. We evaluated the possibility that this difference is a consequence of an E₂ non-genomic signaling pathway present in unmated rats, but not in mated rats. Mating shuts down this E₂ non-genomic signaling pathway up- and downstream of cAMP production. The *Star* level is increased by E₂ in unmated rats, but not in mated rats. This is blocked by the antagonist of estrogen receptor ICI 182 780, the adenylyl cyclase inhibitor SQ 22536, and the catechol-*O*-methyltransferase inhibitor, OR 486. These results indicate that the E₂-induced gene expression profile in the rat oviduct differs before and after mating, and this difference is probably mediated by an E₂ non-genomic signaling pathway operating on gene expression only in unmated rats.

Reproduction (2010) **139** 631–644

Introduction

The oviduct provides an optimal microenvironment for fertilization and early embryo development (Jansen 1984) and delivers the embryo to the uterus at the right time for implantation (Croxatto 1996). Estradiol (E₂) is one of the main regulators of these phenomena modifying expression and secretion of molecules, which assure fertilization and embryo viability (Bui 2002, Bhatt *et al.* 2004). E₂ determines how long oocytes and/or embryos are retained in the oviduct (Forcelledo *et al.* 1986, Croxatto *et al.* 1991), possibly through its action on the activity of muscle and ciliated cells (Villalón & Verdugo 1982, Priyadarsana *et al.* 2004, Ríos *et al.* 2007), which provide the driving force for oviductal egg transport (Croxatto 2002).

E₂ exerts its effects after binding to estrogen receptors (ERs) which belong to the nuclear receptor superfamily (Cheskis *et al.* 2007). The molecular mechanisms

by which E₂ changes the response of target tissues can be through genomic and non-genomic actions. In the genomic actions, estrogens bind to ERs in the nucleus, inducing a conformational change, which results in the regulation of gene transcription. In addition, E₂ is able to activate rapid intracellular signaling pathways (kinase activation, cAMP, and IP₃ increase). Since the latter actions are not blocked by inhibitors of transcription or translation, they have been defined as non-genomic actions (Bjornstrom & Sjoberg 2005).

A single injection of E₂ to unmated or mated rats shortens oviductal transport of eggs from the normal 72–96 h to <24 h (Ortiz *et al.* 1979). In the absence of mating, E₂ uses only intraoviductal non-genomic pathways to accelerate egg transport (Ríos *et al.* 1997, Orihuela & Croxatto 2001, Orihuela *et al.* 2001). This non-genomic pathway involves a previous conversion of E₂ to methoxyestradiols (MEs) mediated by the enzyme catechol-*O*-methyltransferase

(COMT; Parada-Bustamante *et al.* 2007), ER activation (Orihuela *et al.* 2003), and successive activation of the cAMP–protein kinase A (PKA) and phospholipase C–IP₃ signaling cascades (Orihuela *et al.* 2006).

After mating, a profound change occurs in the pathways utilized by E₂ to accelerate egg transport. Instead of using the non-genomic pathway (Orihuela *et al.* 2001), it uses genomic pathways (Ríos *et al.* 1997).

The change in pathways utilized by E₂ to accelerate egg transport, from non-genomic to genomic, has been designated as ‘intracellular path shifting’ (IPS). This IPS caused by mating is a novel example of functional plasticity in well-differentiated cells. Through the effect of protein kinase inhibitors and translation inhibitors on E₂-induced accelerated ovum transport, we demonstrated that among mating-associated signals, the mechanical sensory stimulation of the genital area and the presence of spermatozoa in the uterus are able to elicit IPS (Parada-Bustamante *et al.* 2003, Peñarroja-Matutano *et al.* 2007); however, the mechanisms by which these signals produce this effect have not been elucidated.

IPS involves, at least, shutdown of the E₂ non-genomic signaling pathway, up- and downstream of MEs production, since mating decreases COMT activity and 2ME does not accelerate ovum transport in mated rats (Parada-Bustamante *et al.* 2007); however, the possibility that IPS involves changes or alterations in the genomic signaling pathway, which is used by E₂ to accelerate ovum transport after mating, has not been evaluated. Previous reports indicate that E₂ is able to change the expression of different proteins in the rat oviduct (Mathieu *et al.* 1989, Hermoso *et al.* 1997, Pérez Martínez *et al.* 2006); however, to our knowledge, no studies have been performed to determine whether these responses are different before and after mating. Here, we compared the effect of E₂ on gene expression in the oviduct of unmated and mated rats. We found that E₂ increased the expression of a number of genes common to both conditions, which indicates the presence of a common genomic pathway in unmated and mated rats. Surprisingly, a group of genes decreased its expression in mated rats, but not in unmated rats, and another group of genes increased its expression in unmated rats, but not in mated rats. Therefore, we evaluated whether E₂ non-genomic signaling pathway, present only in unmated rats, is responsible for these differences.

Results

Mating changes the transcript profile induced by E₂ in the oviduct

A microarray analysis using oviductal samples of unmated and mated rats treated s.c. with 10 µg E₂ or vehicle was performed to determine whether E₂ elicits different genomic effects before and after mating.




Figure 1 Changes in oviductal transcriptome induced by E₂ before (unmated) and after (mated) mating. Unmated (N=10) or mated (N=10) rats were injected s.c. with E₂, 10 µg, or vehicle. Three hours later, oviducts were removed to extract their mRNA and perform a microarray analysis. The results obtained represent one affymetrix chip analysis per group performed with a RNA pool obtained from five animals. The number of transcripts that decreased (A) or increased (B) their expression in response to E₂ is shown separately.

Figure 1A and B shows the genes that decreased or increased their expression in response to E₂ respectively. The transcript profile induced by E₂ was very different when this hormone was applied before or after mating. Before mating, E₂ decreased the expression of few genes, but increased the expression of a larger number of genes in comparison to the mated rats. A total of 26 known transcripts decreased their expression in response to E₂ before mating, whereas 49 did so after mating. The comparison of transcripts that decreased their level in both conditions indicated that only 2 decreased their expression exclusively in unmated rats, 25 decreased their expression only in mated rats, and 24 decreased their expression in both conditions (Fig. 1A). The complete lists of the corresponding genes are given in Tables 1–3.

A total of 73 known transcripts increased their expression in response to E₂ before mating, whereas only 42 did so after mating. The comparison of the genes that changed their expression in both conditions indicated that 35 increased their level exclusively in

Table 1 Transcripts that decreased their level in the oviduct of unmated and mated rats 3 h after an estradiol (E₂), 10 µg, treatment.

Probe set ID affymetrix	Accession number	Gene name	Gene symbol	Order of change	
				Unmated rats	Mated rats
1370019_at	NM_031834	Sulfotransferase family 1A, phenol-preferring, member 1	<i>Sult1a1</i>	-2	-2
1368124_at	NM_133578	Dual specificity phosphatase 5	<i>Dusp5</i>	-2	-2
1368894_at	NM_053874	CAP, adenylate cyclase-associated protein, 2 (yeast)	<i>Cap2</i>	-2	-2
1368718_at	NM_017272	Aldehyde dehydrogenase family 1, subfamily A7	<i>Aldh1a7</i>	-2	-2
1371883_at	NM_001007673	Monocyte to macrophage differentiation-associated	<i>Mmd</i>	-2	-2
1368128_at	NM_031598	Phospholipase A2, group IIA (platelets, synovial fluid)	<i>Pla2g2a</i>	-2	-2
1369670_at	NM_031518	Cd200 antigen	<i>Cd200</i>	-2	-2
1368911_at	NM_017099	Potassium inwardly rectifying channel, subfamily J, member 8	<i>Kcnj8</i>	-2	-2.1
1368869_at	NM_001033653	A kinase (PRKA) anchor protein (gravin) 12	<i>Akap12</i>	-2	-2.1
1370310_at	NM_173094	3-Hydroxy-3-methylglutaryl-coenzyme A synthase 2	<i>Hmgcs2</i>	-2	-2.3
1386908_at	NM_022278	Glutaredoxin 1 (thioltransferase)	<i>Glx1</i>	-2	-2.5
1369113_at	NM_019282	Gremlin 1 homolog, cysteine knot superfamily (<i>Xenopus laevis</i>)	<i>Grem1</i>	-2	-2.6
1370228_at	NM_001013110	Transferrin	<i>Tf</i>	-2.1	-2
1387658_at	NM_012947	Eukaryotic elongation factor-2 kinase	<i>Eef2k</i>	-2.1	-2
1368870_at	NM_013060	Inhibitor of DNA binding 2	<i>Id2</i>	-2.1	-2.1
1367631_at	NM_022266	Connective tissue growth factor	<i>Ctgf</i>	-2.1	-3.5
1368154_at	NM_017090	Guanylate cyclase 1, soluble, α3	<i>Gucy1a3</i>	-2.3	-2.5
1367859_at	NM_013174	Transforming growth factor, β3	<i>Tgfb3</i>	-2.3	-2.6
1387850_at	NM_023020	Transmembrane protein with EGF-like and two follistatin-like domains 1	<i>Tmeff2</i>	-2.3	-4.9
1387809_at	NM_053703	MAP kinase kinase 6	<i>Map2k6</i>	-2.5	-2.1
1387074_at	NM_053453	Regulator of G-protein signaling 2	<i>Rgs2</i>	-2.6	-2.6
1368168_at	NM_053380	Solute carrier family 34 (sodium phosphate), member 2	<i>Slc34a2</i>	-2.8	-2
1368025_at	NM_080906	DNA-damage-inducible transcript 4	<i>Ddit4</i>	-2.8	-2.8
1371731_at	NM_001009617	Mesoderm-specific transcript	<i>Mest</i>	-4	-20

unmated rats, only 4 did so exclusively in mated rats, and 38 increased their level in both conditions (Fig. 1B). The complete lists of the corresponding genes are given in Tables 4–6.

These results indicate that mating has a profound effect on the gene expression profile induced by E₂.

E₂ activates a common genomic pathway in unmated and mated rats

Thirty-eight transcripts increased their expression in unmated and mated rats in response to E₂ (Table 4), indicating that a common signaling pathway activated by E₂ is present before and after mating. The transcript encoding creatine kinase brain (*Ckb*) is one of them, and its increase was confirmed by real-time PCR

(unmated rats: 51.6 ± 8.4 ($N=5$) versus 25.7 ± 4.5 ($N=5$) in the control group and mated rats: 39.2 ± 5.6 ($N=5$) versus 20.8 ± 1.9 ($N=5$) in the control group). Since this transcript is regulated by E₂ in other organs by mechanisms that involve estrogen response elements (EREs) and Sp1 sites in its promoter (O'Lone *et al.* 2004), the common signaling pathway would involve, at least, a classical genomic pathway which requires binding of activated ER to these promoter regions.

E₂ increases cAMP level in the rat oviduct before, but not after, mating

E₂ increases cAMP levels in the oviduct of unmated rats through a non-genomic signaling pathway (Orihuela *et al.* 2003). Here, we evaluated whether E₂ is able to

Table 2 Transcripts that decreased their level exclusively in the oviduct of unmated rats 3 h after an estradiol (E₂), 10 µg, treatment.

Probe set ID affymetrix	Accession number	Gene name	Gene symbol	Order of change in unmated rats
1370913_at	NM_138881	Radical S-adenosyl methionine domain containing 2	<i>Rsad2</i>	-2.1
1369153_at	NM_022628	Nephrosis 1 homolog, nephrin (human)	<i>Nphs1</i>	-9.8

Table 3 Transcripts that decreased their level exclusively in the oviduct of mated rats 3 h after an estradiol (E₂), 10 µg, treatment.

Probe set ID affymetrix	Accession number	Gene name	Gene symbol	Order of change in mated rats
1368771_at	NM_134378	Sulfatase 1	<i>Sulf1</i>	-2
1371179_a_at	XM_001077699	Fibroblast growth factor receptor 2 predicted	<i>Fgfr2</i>	-2
1387122_at	NM_012760	Pleiomorphic adenoma gene-like 1	<i>Plagl1</i>	-2
1370363_at	NM_133295	Carboxylesterase 3	<i>Ces3</i>	-2.1
1369526_at	NM_013084	Acyl-coenzyme A dehydrogenase, short/branched chain	<i>Acadsb</i>	-2.3
1368522_at	NM_031340	Timeless homolog (<i>Drosophila</i>)	<i>Timeless</i>	-2.5
1367598_at	NM_012681	Transthyretin	<i>Ttr</i>	-2.5
1387082_at	NM_053348	Fetuin β	<i>Fetub</i>	-2.6
1368543_at	NM_053524	NADPH oxidase 4	<i>Nox4</i>	-2.8
1371083_at	NM_182474	Serine protease inhibitor	<i>LOC299282</i>	-3
1368288_at	NM_012564	Group-specific component	<i>Gc</i>	-3
1368161_a_at	NM_012898	α-2-HS-glycoprotein	<i>Ahsg</i>	-3.5
1368397_at	NM_001007264	UDP-glucuronosyltransferase 2 family, member 5	<i>Ugt2b5</i>	-3.5
1387125_at	NM_053587	S100 calcium-binding protein A9 (calgranulin B)	<i>S100a9</i>	-3.5
1370009_at	NM_012501	Apolipoprotein C-III	<i>Apoc3</i>	-3.7
1370086_at	NM_012559	Fibrinogen γ polypeptide	<i>Fgg</i>	-3.7
1369111_at	NM_012556	Fatty acid-binding protein 1, liver	<i>Fabp1</i>	-5.3
1367896_at	NM_019292	Carbonic anhydrase 3	<i>Ca3</i>	-6.1
1368627_at	NM_031546	Regucalcin	<i>Rgn</i>	-6.5
1367555_at	NM_134326	Albumin	<i>Alb</i>	-11.3
1370396_x_at	NM_001034950	Urinary protein 2	<i>Rup2</i>	-17.1
1368733_at	NM_012883	Sulfotransferase, estrogen preferring	<i>Ste</i>	-22.6
1370593_at	NM_153312	Cytochrome P450, family 3, subfamily a, polypeptide 11	<i>Cyp3a11</i>	-24.3
1370349_a_at	XM_001056871	Similar to urinary protein 3 precursor (RUP-3)	<i>LOC680367</i>	-24.3
1368048_at	NM_012657	Serine (or cysteine) peptidase inhibitor, clade A, member 3K	<i>Serpina3k</i>	-27.9

produce the same response after mating. In unmated rats, E₂ increased cAMP levels almost twofold (2.76 ± 0.52 pmol/oviduct ($N=3$) versus 1.3 ± 0.2 pmol/oviduct ($N=3$) in the control group) (Fig. 2), whereas in mated rats, E₂ did not change it (1.3 ± 0.2 pmol/oviduct ($N=3$) versus 1.2 ± 0.35 ($N=3$) pmol/oviduct in the control group) (Fig. 2), indicating that mating shuts down an E₂ non-genomic signaling pathway in the oviduct, upstream of cAMP generation.

Mating shuts down E₂ non-genomic signaling pathway downstream of cAMP generation

Pharmacological treatments that increase cAMP production in the rat oviduct accelerate ovum transport in unmated rats (Orihuela *et al.* 2003). However, this effect has not been evaluated in mated rats. Twenty-four hours after treatment with forskolin, an average of 40% of oocytes left the oviduct prematurely in unmated rats (6 ± 0.8 ($N=5$) versus 10.2 ± 0.8 ($N=5$) oviductal oocytes in the control group; Fig. 3A), whereas in mated rats, no effect was observed with this dose (9.6 ± 1.2 ($N=5$) versus 10 ± 1.8 ($N=5$) oviductal embryos in the control group). To discard whether this effect of forskolin is only due to a differential activation of adenylyl cyclase isoforms in unmated and mated rats,

the effect of N⁶,2'-O-dibutyryl-adenosine 3',5'-cyclic monophosphate sodium salt (dbcAMP), a cAMP analog, was evaluated. In unmated rats, 50% of the oocytes left the oviduct prematurely after treatment with 400 µg of dbcAMP (7 ± 1.1 ($N=5$) versus 11.6 ± 0.8 ($N=5$) in the control group; Fig. 3B), whereas the same dose had no effect on the number of oviductal embryos in mated rats (8.6 ± 1.0 ($N=5$) versus 9.2 ± 1.2 ($N=5$) in the control group; Fig. 3B), indicating that mating shuts down an E₂ non-genomic signaling pathway used by E₂ to accelerate ovum transport, downstream of cAMP generation too.

E₂-induced non-genomic signaling pathway produces changes in gene expression in unmated rats, and mating shuts down this pathway: the case of STAR expression

E₂ increased the level of 35 transcripts exclusively in unmated rats, suggesting that a signaling pathway activated by E₂ present before, but not after, mating is responsible for this effect. To characterize this response, a temporal course of *Star* expression, one of the transcripts that increased its expression in response to E₂ only in unmated rats, at RNA and protein levels was performed after E₂ treatment.

In unmated rats, E₂ significantly increased *Star* mRNA only 4.5 h after E₂ treatment (61.2 ± 15.2 vs 22.2 ± 6.1

Table 4 Transcripts that increased their level in the oviduct of unmated and mated rats 3 h after an estradiol (E₂), 10 µg, treatment.

Probe set ID affymetrix	Accession number	Gene name	Gene symbol	Order of change	
				Unmated rats	Mated rats
1367894_at	XM_001066529	Similar to insulin-induced gene 1 protein (INSIG-1)	<i>LOC688922</i>	+2	+2
1368685_at	NM_031022	Chondroitin sulfate proteoglycan 4	<i>Cspg4</i>	+2	+2
1387184_at	NM_024355	Axin2	<i>Axin2</i>	+2	+2.1
1368542_at	NM_053583	Zinc finger protein 423	<i>Zfp423</i>	+2	+2
1368272_at	NM_012571	Glutamate oxaloacetate transaminase 1	<i>Got1</i>	+2	+3
1367802_at	NM_019232	Serum/glucocorticoid-regulated kinase	<i>Sgk</i>	+2	+2.3
1370032_at	NM_021594	Solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulator 18	<i>Slc9a3r1</i>	+2	+2.8
1368465_at	NM_012892	Amiloride-sensitive cation channel 1, neuronal (degenerin)	<i>Accn1</i>	+2	+2
1387908_at	XM_001077321	RAS, dexamethasone-induced 1	<i>Rasd1</i>	+2.1	+2
1387391_at	NM_080782	Cyclin-dependent kinase inhibitor 1A	<i>Cdkn1a</i>	+2.1	+4.9
1367740_at	M14400	Creatine kinase, brain	<i>Ckb</i>	+2.1	+2
1369958_at	NM_022542	Ras homolog gene family, member B	<i>Rhob</i>	+2.1	+2
1370074_at	XM_001055943	Tensin	<i>Tns</i>	+2.1	+2
1370884_at	M36410	Sepiapterin reductase	<i>Spr</i>	+2.1	+2.1
1387091_at	NM_017226	Peptidyl arginine deiminase, type II	<i>Padi2</i>	+2.1	+2
1370379_at	NM_138836	Protease, serine, 8 (prostasin)	<i>Prss8</i>	+2.1	+2
1388133_at	XM_345860	Cold shock domain containing C2, RNA binding	<i>Csdc2</i>	+2.1	+2
1368231_at	NM_017064	Stat5A	<i>Stat5a</i>	+2.1	+2
1387332_at	NM_022275	Neuromedin U receptor 2	<i>Nmur2</i>	+2.1	+2.1
1387010_s_at	NM_017288	Sodium channel, voltage-gated, type I, β	<i>Scn1b</i>	+2.1	+3.7
1368826_at	NM_012531	Catecholamine-O-methyltransferase	<i>Comt</i>	+2.1	+2
1368223_at	NM_024400	ADAM metalloproteinase with thrombospondin type 1 motif, 1	<i>Adams1</i>	+2.3	+2
1369520_a_at	NM_017253	Branched chain aminotransferase 1, cytosolic	<i>Bcat1</i>	+2.3	+2.3
1387099_at	NM_053838	Natriuretic peptide receptor 2	<i>Npr2</i>	+2.3	+2
1370173_at	NM_017051	Superoxide dismutase 2, mitochondrial	<i>Sod2</i>	+2.3	+2.1
1387563_at	NM_022847	Progesterone receptor	<i>Pgr</i>	+2.3	+3.2
1387737_at	NM_134351	Methionine adenosyltransferase II	<i>Mat2</i>	+2.3	+2
1387260_at	NM_053713	Kruppel-like factor 4 (gut)	<i>Klf4</i>	+2.5	+2
1370333_a_at	NM_178866	Insulin-like growth factor 1	<i>Igf1</i>	+2.5	+2
1370225_at	NM_053699	Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 4	<i>Cited4</i>	+2.6	+2.3
1367725_at	NM_022602	Serine/threonine-protein kinase pim-3	<i>Pim3</i>	+2.6	+2
1368283_at	NM_133606	Enoyl-coenzyme A, hydratase/3-hydroxyacyl coenzyme A dehydrogenase	<i>Ehhadh</i>	+2.6	+2
1369443_at	NM_133569	Angiopoietin-like 2	<i>Angptl2</i>	+3	+2
1370708_a_at	NM_138547	Aldo-keto reductase family 1, member C14	<i>Akr1c14</i>	+3	+2.1
1369798_at	NM_012507	ATPase, Na ⁺ /K ⁺ transporting, β2 polypeptide	<i>Atp1b2</i>	+4	+2
1368148_at	NM_012610	Nerve growth factor receptor (TNFR superfamily, member 16)	<i>Ngfr</i>	+5.3	+2
1370517_at	NM_153735	Neuronal pentraxin 1	<i>Nptx1</i>	+9.2	+3
1368102_at	NM_017081	Hydroxysteroid 11β dehydrogenase 2	<i>Hsd11b2</i>	+9.2	+3.5

group vehicle, Fig. 4, unmated), whereas its protein level was increased at all times studied, with maximal increases being reached 4.5 and 6 h after E₂ treatment (4.0 ± 1.0 and 4.5 ± 1.2 times respectively, Fig. 5, unmated). In mated rats, E₂ did not change mRNA (Fig. 4, mated) and protein (Fig. 5, mated) levels of STAR at any time studied.

Since mating shuts down E₂-induced non-genomic signaling pathway up- and downstream of cAMP

generation, the possibility that this pathway is responsible for transcript level changes induced by E₂ observed only in unmated rats was evaluated. For this, the E₂-induced *Star* expression increase in unmated rats was studied by blocking the ER and cAMP production, which are the two key components that mediate E₂ non-genomic signaling pathway (Orihuela *et al.* 2003). ICI 182 780, an ER antagonist, and SQ 22536, an adenylyl cyclase inhibitor, or their vehicles were administered

Table 5 Transcripts that increased their level exclusively in the oviduct of unmated rats 3 h after an estradiol (E₂), 10 µg, treatment.

Probe set ID affymetrix	Accession number	Gene name	Gene symbol	Order of change in unmated rats
1369554_at	NM_053553	Synaptogyrin 2	<i>Syngn2</i>	+2
1387629_at	NM_022261	B-box and SPRY domain containing	<i>Bspry</i>	+2
1370153_at	NM_019216	Growth differentiation factor 15	<i>Gdf15</i>	+2
1371059_at	NM_019264	Protein kinase, cAMP-dependent, regulatory, type 2 α	<i>Prkar2a</i>	+2
1369177_at	NM_053735	Phosphatidylinositol 4-kinase type 2 α	<i>Pi4k2a</i>	+2
1370212_at	NM_053310	Homer homolog 3 (<i>Drosophila</i>)	<i>Homer3</i>	+2
1368845_at	NM_024000	caM kinase-like vesicle-associated	<i>Camkv</i>	+2
1368147_at	NM_053769	Dual specificity phosphatase 1	<i>Dusp1</i>	+2
1368449_at	NM_133567	Centaurin α 1	<i>Centa1</i>	+2
1387894_at	NM_144730	GATA binding protein 4	<i>Gata4</i>	+2
1369012_at	NM_017128	Inhibin β -A	<i>Inhba</i>	+2
1387353_at	NM_017093	Thymoma viral proto-oncogene 2	<i>Akt2</i>	+2
1387876_at	NM_022380	Stat5B	<i>Stat5b</i>	+2.1
1386909_a_at	NM_031353	Voltage-dependent anion channel 1	<i>Vdac1</i>	+2.1
1369650_at	NM_053306	p21 (CDKN1A)-activated kinase 2	<i>Pak2</i>	+2.1
1368931_at	NM_031238	SH3-domain GRB2-like 3	<i>Sh3gl3</i>	+2.1
1387133_at	NM_053988	Calbindin 2	<i>Calb2</i>	+2.3
1387844_at	NM_032613	LIM and SH3 protein 1	<i>Lasp1</i>	+2.3
1370780_at	NM_145094	RAB31, member RAS oncogene family	<i>Rab31</i>	+2.3
1368884_at	NM_022587	Ectonucleoside triphosphate diphosphohydrolase 1	<i>Entpd1</i>	+2.3
1369640_at	NM_012567	Gap junction membrane channel protein α 1	<i>Gja1</i>	+2.5
1369278_at	NM_031034	Guanine nucleotide binding protein, α 12	<i>Gna12</i>	+2.5
1383096_at	XM_343513	Amyloid β (A4) precursor-like protein 2	<i>Aplp2</i>	+2.5
1387707_at	NM_017102	Solute carrier family 2 (facilitated glucose transporter), member 3	<i>Slc2a3</i>	+2.5
1387810_at	NM_057152	Kelch-like ECH-associated protein 1	<i>Keap1</i>	+2.5
1370488_a_at	NM_019140	Protein tyrosine phosphatase, receptor type, D	<i>Ptprd</i>	+2.6
1387420_at	NM_031818	Chloride intracellular channel 4	<i>Clic4</i>	+2.8
1387858_at	NM_080907	Protein phosphatase 4, regulatory subunit 1	<i>Ppp4r1</i>	+2.8
1370344_at	NM_153629	Heat shock protein 4	<i>Hspa4</i>	+3
1368406_at	NM_031558	Star	<i>Star</i>	+3.2
1370714_a_at	NM_147205	β -Galactoside- α -2,6 sialyltransferase 1	<i>St6gal1</i>	+3.2
1369703_at	NM_023090	Endothelial PAS domain protein 1	<i>Epas1</i>	+3.2
1368897_at	NM_030858	MAD homolog 7 (<i>Drosophila</i>)	<i>Smad7</i>	+3.5
1387833_at	NM_130424	Transmembrane protease, serine 2	<i>Tmprss2</i>	+6.5
1388249_at	XM_001079347	Rap guanine nucleotide exchange factor (GEF) 1	<i>Rapgef1</i>	+10.6

intrabursally (i.b.) in unmated rats following a s.c. injection with E₂ or its vehicle. In rats receiving local treatment with vehicle, E₂ increased STAR levels (ICI experiment: 2.16 ± 0.17 ($N=3$), Fig. 6A, and SQ 22536 experiment: 1.81 ± 0.01 ($N=3$), Fig. 6B). ICI 182 780 and SQ 22536 completely blocked the increase of STAR induced by E₂ (0.65 ± 0.07 ($N=3$) and 1.12 ± 0.16 ($N=3$), Fig. 6A and B respectively), whereas these drugs when applied alone did not affect STAR expression by themselves (0.8 ± 0.14 ($N=3$) and 1.09 ± 0.11 ($N=3$), Fig. 6A and B respectively). These results

indicate that increased STAR expression induced by E₂ in unmated rats requires an active ER and an increase of cAMP levels.

STAR immunoreactivity was detected only in epithelial cells of the ampulla and isthmus (Fig. 7), indicating that at least part of the pathway utilized by E₂ to increase STAR expression is present in this cell phenotype.

Altogether these data indicate that some, if not all, genes that increase their expression in the oviductal epithelium in response to E₂ exclusively in unmated rats do so, at least in part, in response to the activation of a

Table 6 Transcripts that increased their level exclusively in the oviduct of mated rats 3 h after an estradiol (E₂), 10 µg, treatment.

Probe set ID affymetrix	Accession number	Gene name	Gene symbol	Order of change in mated rats
1368339_at	NM_012521	S100 calcium-binding protein G	<i>S100g</i>	+2.5
1368342_at	NM_031544	Adenosine monophosphate deaminase 3	<i>Ampd3</i>	+2.8
1368290_at	NM_031327	Cysteine-rich protein 61	<i>Cyr61</i>	+4
1371193_at	XM_001065494	Tumor necrosis factor α -induced protein 6	<i>Tnfaip6</i>	+10




Figure 2 Effect of estradiol on oviductal cAMP level before (unmated) and after (mated) mating. cAMP level in the oviduct of unmated and mated rats ($N=3$ animals by treatment) 3 h after a s.c. injection of E₂, 10 μg , or vehicle. Numbers inside the bars indicate the number of animals used. Means with different letters are significantly different from each other ($P<0.05$).

local non-genomic signaling pathway that increases cAMP levels. Mating shuts down this pathway, preventing E₂ from increasing the expression of these genes in mated rats.

E₂ regulated STAR expression, COMT requirement, and effect of 2ME

Local formation of MEs mediated by COMT is essential for the intraoviductal non-genomic pathway utilized by E₂ to accelerate egg transport in unmated rats (Parada-Bustamante *et al.* 2007). Here, we evaluated whether OR 486, a COMT inhibitor, blocks the E₂-induced increase in STAR levels. OR 486 or its vehicle was administered i.b. in unmated rats after a s.c. injection of E₂ or its vehicle. E₂ increased STAR levels significantly (2.3 ± 0.6 times over control, Fig. 8), and OR 486 completely blocked this effect (1.1 ± 0.2 times over control, Fig. 8), whereas OR 486 alone had no effect (1.2 ± 0.3 times over control). Moreover, STAR levels increased significantly 1.5 and 3 h after 2ME treatment (2.4 ± 0.6 and 1.6 ± 0.3 times over control respectively, Fig. 9), indicating that local conversion of E₂ to ME is an essential component of the non-genomic signaling pathway of E₂ that regulates STAR expression in the oviduct of unmated rats.

Discussion

Mating has a profound impact on oviductal physiology, changing the pathway by which E₂ accelerates ovum transport from non-genomic to genomic (IPS). In this study, the genomic pathways that respond to E₂ before and after mating were explored. We found that the E₂-induced gene expression profile in the rat oviduct is different depending on whether the animal has mated or not. More genes were increased and fewer genes were

decreased by E₂ before mating than after mating. Since more information about the mechanisms by which E₂ increases gene expression is present in the literature, our analysis was focused on the group of genes that were increased by E₂. The fact that 38 genes are increased in unmated and mated rats indicates that a common pathway that is able to increase gene expression is present before and after mating. According to the literature, some of these genes are regulated by E₂ in other rat organs such as angiotensin 2 in the heart, kidney, and lung (Ye *et al.* 2004), hydroxysteroid 11 β dehydrogenase 2 in the kidney (Gómez-Sánchez *et al.* 2003), insulin-like growth factor 1 (*Igf1*) in the oviduct (Carlsson *et al.* 1993), progesterone receptor in the lung (González-Arenas *et al.* 2003), and *Ckb* in pituitary gland (Blake *et al.* 2005). Functional analysis of progesterone receptor promoter, which is one among this group of genes, indicated that it contained imperfect EREs, which are necessary for increasing its levels by E₂; moreover, *Ckb* promoter contains EREs and GC-rich sites, which are used by E₂, bound to ER, to increase its levels (Scott *et al.* 2003). In this work, we confirmed that E₂ increased *Ckb* mRNA levels in unmated and mated rat




Figure 3 Forskolin and dibutylryl-cAMP accelerate ovum transport in unmated rats, but not in mated rats. Number of oocytes or embryos recovered from the oviduct of unmated and mated rats ($N=5$ animals by treatment) 24 h after being injected with forskolin, 20 μg , or vehicle i.b. (A) or with db-cAMP, 4 or 400 μg , or vehicle i.b. (B). Numbers inside the bars indicate the number of animals used. Means with different letters are significantly different from each other ($P<0.05$).




Figure 4 Estradiol increases oviductal *Star* mRNA levels in unmated rats, but not in mated rats. *Star* mRNA relative expression obtained through real-time PCR from cDNA of oviductal samples ($N=5$ animals by treatment) taken from unmated and mated rats 3, 4.5, 6, 12, and 24 h after a s.c. injection of E₂, 10 μ g, or vehicle. The values were normalized to *Gapdh*. Each bar represents the mean value obtained from five animals. The asterisk indicates a statistically significant difference between the two bars at 4.5 h ($P<0.05$).

oviduct, indicating that the common pathway is, at least, a classical genomic pathway which is dependent on direct or indirect ER–DNA interaction.

E₂ increased the expression of some transcripts exclusively in unmated rats, but not in mated rats, indicating that a pathway that is able to increase the expression of these transcripts is present in unmated rats, whereas mating shut it down by unrecognized mechanisms. Previously, we had determined that in IPS, the E₂ non-genomic pathway is shut down up- and downstream of 2ME (Parada-Bustamante *et al.* 2007); here, we show that mating shuts down the non-genomic pathway up- and downstream of cAMP production. The fact that the E₂ non-genomic signaling pathway is shut down due to mating, and that some genes, such as inhibin β -A (Ardekani *et al.* 1998), *Smad7* (Bilezikjian *et al.* 2001), gap junction protein, α 1 (Abudara *et al.* 1999), and *Star* (Stocco *et al.* 2005), which increased their expression only in unmated rats are increased by

cAMP and PKA activation in other rat tissues, indicates that in unmated rats, E₂ increases the expression of these genes activating a non-genomic pathway. After mating, this pathway is shut down and then E₂ is unable to increase the expression level of this group of genes. In accordance with this idea, in unmated rats, *Star* mRNA levels are increased 4.5 h after E₂ treatment, whereas in mated rats, *Star* mRNA levels did not change at any time point studied. Furthermore, the fact that increases of STAR protein induced by E₂ require ER activation and cAMP production suggests that this effect is dependent on the intraoviductal E₂ non-genomic signaling. Since ICI 182 780 alone did not affect STAR expression, it is




Figure 5 Estradiol increases oviductal STAR protein levels in unmated rats, but not in mated rats. Densitometric analysis of western blots to detect STAR protein in oviductal samples (20 μ g, $N=3$ animals by treatment) taken 3, 4.5, 6, and 7.5 h after a s.c. injection of E₂, 10 μ g, or vehicle. The values were normalized to α -tubulin. A representative western blot obtained in this experiment is shown. Numbers inside the bars indicate the number of animals used. Means with different letters are significantly different from each other ($P<0.05$).




Figure 6 Increased expression of STAR levels in the oviduct in response to E₂ requires estrogen receptor and cAMP. Densitometric analysis of western blots to detect STAR protein in oviductal samples (20 µg, N=3 animals by treatment) of un-mated rats 4.5 h after injecting E₂, 10 µg, or vehicle s.c. and concomitantly the estrogen receptor antagonist, ICI 182 780 (A), the adenylyl cyclase inhibitor, SQ 22536 (B), or their vehicle given i.b. The values were normalized to α -tubulin. A representative western blot is shown. Numbers inside the bars indicate the number of animals used. Means with different letters are significantly different from each other ($P < 0.05$).

probable that other endogenous signaling pathways are acting to state basal STAR levels in the rat oviduct. In accordance with this idea, previous reports have determined that STAR expression is regulated by several signaling pathways and factors, such as IGF1, arachidonic acid, protein kinase C, and MAP kinases (reviewed in Stocco *et al.* (2005)).

The fact that *Star* mRNA levels were increased only in one of the points studied is in agreement with our previous results, which indicated that cAMP levels are increased 3 h after E₂ treatment, whereas its levels returned to basal levels at 6 h (P Orihuela 2006, unpublished observations). STAR protein levels were

increased slightly, but significantly 3 h after E₂ treatment, even though its mRNA levels were increased from 4.5 h onwards after treatment, probably because cAMP–PKA pathway activation increases STAR protein stability (Clark *et al.* 2001). This also indicates that STAR protein levels are maintained elevated until 7.5 h after E₂ treatment. In mated rats, E₂ did not change STAR protein levels at any time studied. These results indicate that E₂ increases STAR protein levels in unmated rats, activating a non-genomic pathway that requires E₂ receptor and adenylyl cyclase.

This shutdown of E₂ non-genomic signaling pathway in mated rats could be explained by differences in E₂ levels or changes in the expression or location of ER induced by mating in the rat oviduct. However, previous reports have shown that plasma E₂ levels and oviductal ESR1 (ER- α) and ESR2 (ER- β) mRNA and protein levels did not change in the initial hours after mating (Smith *et al.* 1975, Orihuela *et al.* 2004). Another possibility could be the interactions among ERs and proteins that mediate E₂ non-genomic signaling pathways in unmated rats, which do not occur in mated rats. For example, it has been found that PELP-1/MNAR is an ER-interacting protein (Brann *et al.* 2008) which is required for ESR1 interaction with p60 (SRC), which leads to the activation of SRC/MAPK pathway (reviewed in Cheskis *et al.* (2008)); however, this possible differential interaction in the rat oviduct was not evaluated in this work.

E₂ increased STAR levels only in the epithelial cells, indicating that E₂-activated non-genomic pathway is present only in these cells. This is in agreement with our previous results using primary cell cultures from rat oviduct, where E₂ increased cAMP levels in epithelial cells, but not in muscle cells (P Orihuela 2007, unpublished observations); however, we do not discount that other components of the non-genomic pathway are present in other cell types, since IP₃ levels are increased 1 and 6 h after E₂ treatment in the whole rat oviduct (Orihuela *et al.* 2006).

cAMP regulates STAR expression in other tissues and organs through PKA activation and subsequent transcription factor phosphorylation such as GATA4 and CREB (Stocco *et al.* 2005). The effectors that are downstream of cAMP production, responsible for E₂-induced STAR increases in the rat oviduct, were not studied in this work. To our knowledge, this is the first time that STAR expression is reported in the oviduct, and that its expression is increased by E₂; a previous work reported that STAR expression is decreased in interstitial and theca ovarian cells in newborn rats treated with E₂ benzoate (Ikeda *et al.* 2001). STAR is a key protein in steroidogenesis, because it mediates cholesterol entry from external to internal mitochondrial membranes (Stocco & Clark 1996). The function of STAR in the oviduct was not explored in this work; however, since E₂-induced acceleration transport is not blocked by transcription and translation inhibitors in unmated rats,




Figure 7 STAR is expressed in epithelial cells in unmatred rat oviduct. Representative photomicrographs obtained from unmatred rat oviducts 4.5 h after injecting E_2 , 10 μ g, s.c. ($N=3$) to detect STAR expression (green). Nuclei were stained with propidium iodide (red). Arrows point to immunoreactivity obtained only in the epithelial cells. The specificity of immunoreactivity was assessed by incubating samples with preimmune serum.

the increase of its expression would not mediate this phenomenon.

The E_2 non-genomic pathway in the oviduct involves conversion of E_2 to MEs mediated by COMT (Parada-Bustamante *et al.* 2007). In accordance with this, a COMT inhibitor blocked E_2 -induced STAR increased expression and 2ME increased STAR protein expression. The fact that STAR levels were increased only 3 h after 2ME treatment, whereas they were increased 3, 4.5, 6, and 7.5 h after E_2 treatment can be explained by a fast 2ME metabolism. cAMP levels are increased in the rat oviduct only 3 h after E_2 treatment (Orihuela *et al.* 2006); this latency time can represent the time required by E_2 to be metabolized to 2ME. We hypothesize that 2ME treatment would increase cAMP faster than E_2 , but this was not tested in this work.

The physiological relevance of preventing increased expression of a group of genes by E_2 in the rat oviduct by mating-associated signals was not explored in this work. We postulate that failure of the mechanism that is responsible for the shutdown of this pathway would affect normal embryo development. In order to corroborate this idea, it is necessary to prevent IPS and then to determine whether in this condition there are alterations in reproductive phenomena such as embryo development and implantation; however, this was not explored in this work.

In summary, these results indicate that E_2 -induced gene expression profile in the rat oviduct differs before and after mating, and this difference is possibly mediated by the effects of E_2 non-genomic signaling pathway on gene expression operating only in unmatred rats; the early events induced by mating responsible for this phenomenon are still unknown.

Materials and Methods

Animals

Locally bred Sprague–Dawley rats were used. The animals were kept under controlled temperature (21–24 °C), and lights were kept on from 0700 to 2100 h. Water and pelleted rat chow were supplied *ad libitum*. Females weighing 200–220 g were selected from those that had at least two regular cycles of 4 days immediately before the experiments were started.

Daily vaginal smears, which were taken between 0800 and 0900 h, were used to verify cycle regularity (Turner 1961). To obtain unmatred and mated rats, females in the evening of proestrus were either kept isolated or caged with fertile males. The next morning, isolated rats that presented cornified cells in the vaginal smear, a cell phenotype associated with ovulation (estrus day), were designated as unmatred rats, and those caged with fertile males that presented cornified cells and spermatozoa in the vaginal smear were designated as mated rats. The care and manipulation of the animals were carried out in accordance with the ethical guidelines of Pontificia Universidad Católica de Chile and Universidad de Santiago de Chile.

Treatments

All treatments described below were administered at 1200 h in unmatred and mated rats.




Figure 8 OR 486 blocks increased expression of STAR levels in the oviduct in response to E_2 in unmatred rats. Densitometric analysis of western blots to detect STAR protein in oviductal samples (20 μ g, $N=3$ animals by treatment) of unmatred rats 4.5 h after injecting E_2 , 10 μ g, or vehicle s.c. and concomitantly the COMT inhibitor, OR 486, or vehicle given i.b. The values were normalized to α -tubulin. A representative western blot is shown. Numbers inside the bars indicate the number of animals used. Means with different letters are significantly different from each other ($P<0.05$).




Figure 9 2-Methoxyestradiol increases oviductal STAR levels in unimpaired rats. Densitometric analysis of western blots to detect STAR protein in oviductal samples (20 µg, N=3 animals by treatment) from unimpaired rats taken 1.5 and 3 h after a s.c. injection of 2-methoxyestradiol, 100 µg, or vehicle. The values were normalized to α -tubulin. A representative western blot is shown. Numbers inside the bars indicate the number of animals used. Means with different letters are significantly different from each other ($P < 0.05$).

Systemic administration of E₂ or 2ME

Unmated and mated rats were injected s.c. with 10 µg of E₂ as a single dose in an injection volume of 0.1 ml of propylene glycol. Other rats received a single s.c. injection of 100 µg of 2ME (Steraloids, Newport, RI, USA) in an injection volume of 0.1 ml of propylene glycol. Control rats received propylene glycol alone.

Local administration of drugs

Unmated and mated rats were injected in the ovarian bursa (i.b.) with one of the drugs described below. Control rats received the appropriate vehicle only. Forskolin (7 β -acetoxy-8, 13-epoxy-1,6 β ,9-trihydroxy-labd-14-ene-11-one; Sigma Chemical) and dbcAMP (Sigma Chemical) were used to determine whether E₂ non-genomic signaling pathway is functional downstream of cAMP generation in mated rats using acceleration of ovum transport as end point response. Forskolin, 20 µg, dissolved in 4 µl of 25% ethanol and dbcAMP, 400 µg, dissolved in saline were injected i.b. in unimpaired and mated rats. Previously, we had determined that this dose of forskolin applied i.b. accelerates ovum transport in unimpaired rats (Orihuela *et al.* 2003). ICI 182 780 (kindly donated by W Elger, Entech, Jena, Germany) was used to block the ER, and to determine whether ER is required for increased expression of STAR induced by E₂. ICI 182 780, 25 µg, dissolved in 4 µl of 25% DMSO was injected i.b. in unimpaired rats immediately before injecting 10 µg of E₂ s.c. This dose of ICI 182 780 completely blocks the effect of E₂ on egg transport (Orihuela *et al.* 2003). SQ 22536, an adenylyl cyclase inhibitor, was used to determine whether the E₂ non-genomic signaling pathway is responsible for increasing STAR protein level in unimpaired rats. SQ 22536,

30 µg, dissolved in saline was injected i.b. in unimpaired rats immediately before injecting 10 µg of E₂ s.c. OR 486 (Tocris, Langford, UK) was used to inhibit COMT activity in the oviduct, and to determine whether local conversion of E₂ to MEs is important to increase STAR protein level in unimpaired rats. OR 486, 125 µg, dissolved in 4 µl of 25% ethanol was injected i.b. in unimpaired rats immediately before injecting 10 µg of E₂ s.c.

Once oviducts were removed, all manipulations were done on an ice-cooled plate.

Measurement of cAMP levels

Three hours after E₂ or vehicle injection, unimpaired (N=6) or mated (N=6) rats were killed and their oviducts were flushed individually. Oviducts in groups of two (one rat) were homogenized in 0.5 ml of ice-cold 10% (v/v) trichloroacetic acid and centrifuged for 15 min at 2200 g at 4 °C. The pellet was discarded, and the supernatant was washed four times with five volumes of water-saturated diethyl ether. The upper layer was discarded after each wash. Following the last wash, the aqueous extract was dried under a stream of nitrogen at 60 °C. Levels of cAMP in dried extracts were determined using Biotrak cAMP enzyme immunoassay system (catalog no. RPN 225; Amersham Pharmacia Biotech, Piscataway, NJ, USA). This kit is based on the competition between unlabeled cAMP and a fixed quantity of peroxidase-labeled cAMP for a limited number of binding sites on a cAMP-specific antibody. This allows for the construction of a standard curve and the measurement of cAMP levels in unknown samples. Color was developed with 3,3',5,5'-tetramethylbenzidine/hydrogen peroxide as a substrate. Optical density was read at 630 nm with a microplate reader (BIO-TEK Instruments, Winooski, VT, USA). Previously, we had determined that E₂ increases cAMP levels at this time in the oviduct of unimpaired rats through a non-genomic signaling pathway (Orihuela *et al.* 2006; and P Orihuela 2006, unpublished observations).

Animal surgery

Intrabursal administration, which minimizes the dose needed to affect the oviduct, avoiding systemic effects, was performed on unimpaired and mated rats at 1200 h as described by Orihuela & Croxatto (2001). Briefly, the oviduct and ovary were exposed through flank incisions made under anesthesia, and using a surgical microscope (OPMI 6-SDFC; Zeiss, Oberkochen, Germany), the drugs or vehicle alone was injected into the periovarian sac using a Hamilton syringe (Hamilton Co., Reno, NV, USA), and the injection site in the bursa was immediately sealed with an electric coagulator (Codman CMC-1; Codman and Shurleff Inc., Randolph, MA, USA). The organs were returned to the peritoneal cavity, and the muscles and skin were sutured. Since ovulation was completed at this time point, this treatment did not affect the number of oocytes that ovulated. Furthermore, we had previously demonstrated that drugs that are administered i.b. act locally on the oviduct (Orihuela & Croxatto 2001, Orihuela *et al.* 2006).

Assessment of egg transport

Animals were killed 24 h after treatment, and their oviducts were flushed individually with saline. Each flushing was examined under low-power magnification (25×). The number of eggs in both oviducts was recorded as a single datum. We had previously determined that the recovery of eggs using this method is close to 100% by comparing the average number of eggs obtained from oviducts with this technique with the number of implanted embryos on day 12 of pregnancy (Ortiz *et al.* 1979). Attempts to recover eggs from the uterus and vagina with or without placing ligatures in the uterine horns have shown that the reduction in the number of oviductal oocytes following treatment with E₂ corresponds to premature transport to the uterus (Ortiz *et al.* 1979). Thus, we refer to this phenomenon as E₂-induced acceleration of oviductal transport.

Oviduct collection and microarray analysis

Unmated (*N*=10) and mated (*N*=10) rats were injected s.c. with 10 µg of E₂ (*N*=5) or vehicle (*N*=5) at noon, and 3 h later, they were killed and their oviducts were collected and flushed. Total oviductal RNA was isolated from each rat using Trizol reagent (Invitrogen), and equivalent quantities were mixed to generate four pools: unmated group treated with E₂ (unmated group E₂), unmated group treated with vehicle (unmated group vehicle), mated group treated with E₂ (mated group E₂), and mated group treated with vehicle (mated group vehicle). RNA probes prepared from each group were hybridized by Genome Explorations Inc. to the Rat Genome 230 2.0 chips (Affymetrix GeneChip System; Affymetrix, Santa Clara, CA, USA) according to the manufacturer's instructions. The transcriptome profile of the genes that were increased or decreased by E₂ in the unmated group was compared to that of those that were increased or decreased by E₂ in the mated group.

Real-time PCR

In order to determine the response of *Star* and *Ckb* RNA expression in the oviduct to E₂, unmated and mated rats were injected s.c. with 10 µg of E₂ or vehicle, and they were killed at various time intervals and their oviducts were collected and flushed. Total oviductal RNA was isolated from each rat using RNAsolv (Omega Bio-Tek, Norcross, GA, USA), and 1 µg of total RNA of each sample (two oviducts from one rat) was treated with DNase I (amplification grade; Invitrogen). The single-strand cDNA was synthesized by reverse transcription using the Superscript III Reverse Transcriptase First Strand System for RT-PCR (Invitrogen) according to the manufacturer's protocol. The Light Cycler instrument (Roche Diagnostics) was used to quantify the relative gene expression of *Star* and *Ckb* in the oviducts of unmated and mated rats treated with E₂ or vehicle; *Gapdh* was chosen as the housekeeping gene for loading control. The SYBR Green I double-strand DNA binding dye from QuantiTec Real-Time RT-PCR kit (Qiagen) was the reagent of choice for these assays. The following primers were used: for *Star*, sense 5'-CTG CTA GAC CAG CCC ATG GAC-3' and antisense 5'-TGA TTT CCT TGA CAT TTG GGT TCC-3'; for *Ckb*, sense 5'-AAG CTG GCA GTA GAA GCC CT-3' and

antisense 5'-TTG TCG AAG AGG AAG TGG TC-3'; and for *Gapdh*, sense 5'-ACC ACA GTC CAT GCC ATC AC-3' and antisense 5'-TCC ACC ACC CTG TTG CTG TA-3'. All real-time PCR assays were performed in duplicate. The thermal cycling conditions included an initial activation step at 95 °C for 25 min, followed by 40 cycles of 95 °C for 15 s, 59 °C for 30 s, and 72 °C for 30 s, with an ultimate melting cycle (95–60 °C). In order to verify the specificity of each product, amplified products were subjected to melting curve analysis as well as to electrophoresis, and product sequencing was performed using an ABI Prism 310 sequencer. The expression of *Star* was determined using the equation: $Y = 2^{-\Delta C_p}$, where *Y* is the relative expression, *C_p* (crossing point) is the cycle in the amplification reaction in which fluorescence begins to expand exponentially above the background baseline, and $-\Delta C_p$ is the result of subtracting the *C_p* value of *Star* from the *C_p* value of *Gapdh* for each sample. To simplify the presentation of the data, the relative expression values were multiplied by 10³ (Livak & Schmittgen 2001).

Immunoblotting

Oviducts obtained from unmated and mated rats (*N*=3 animals for each described experiment) were flushed, and their total proteins were isolated as described by Irueta *et al.* (2003). Briefly, oviducts were lysed in lysis buffer (20/ml³ Tris-HCl, pH 8.0, 137/ml³ NaCl, 1% Nonidet P-40, and 10% glycerol) supplemented with a protease inhibitor cocktail (Complete; Roche). The lysate was centrifuged at 4 °C for 10 min at 10 000 *g*, and the pellet was discarded. Protein concentrations in the supernatant were measured by the Bradford assay (Bio-Rad). After boiling for 5 min, proteins (20 µg) were separated on 15% SDS-PAGE slab gels in a Mini PROTEAN electrophoretic chamber (Bio-Rad). Proteins resolved in the gels were electroblotted onto nitrocellulose membranes (Bio-Rad). The membranes were blocked for 3 h in TTBS (100/ml³ Tris-HCl (pH 7.5), 150/ml³ NaCl, 0.05% (v/v) Tween-20) that contained 5% non-fat dry milk, and were incubated overnight with rabbit anti-STAR (kindly donated by Dr Douglas Stocco, Texas University) or mouse anti- α -tubulin antibody (T5168) Sigma Chemical) at 1:1500 or 1:5000 dilution respectively. The blots were rinsed five times for 5 min each in TBS (100/ml³ Tris-HCl, pH 7.5, and 150/ml³ NaCl) and were incubated for 1 h in TTBS that contained HRP-conjugated goat anti-rabbit or anti-mouse IgG (1:5000 dilution; Chemicon, Billerica, MA, USA). HRP activity was detected by ECL using Western Lightning Chemiluminescence Reagent Plus (PerkinElmer Life Sciences, Boston, MA, USA). Oviductal samples without anti-STAR or anti-tubulin antibody were included as negative controls.

Immunohistochemistry

Oviducts from unmated rats treated s.c. with E₂ (*N*=3 animals) and sacrificed 4.5 h later were fixed in cold 4% paraformaldehyde in PBS, pH 7.4–7.6, for 2 h, and then a sequential transfer to 10% w/v sucrose in PBS for 60 min at 4 °C and 30% w/v sucrose in PBS at 4 °C overnight was done.

Cryostat sections, 4–6 µm thick, were placed onto gelatin-coated slides and were blocked with 1% PBS-BSA for 120 min,

and then incubated with 1:100 anti-STAR antibody in 1% PBS–BSA in a humidified chamber overnight. Three PBS rinses were followed by 60-min incubation at room temperature with secondary antibody biotin-conjugated anti-rabbit IgG (Biosource, Nivelles, Belgium) diluted in 1% PBS–BSA. After three PBS rinses, the slides were incubated with avidin–FITC (Sigma) that was diluted 1:5000 for 60 min at room temperature. Samples were subsequently washed with PBS, counterstained with 1 µg/ml propidium iodide, and mounted in DABCO (Sigma). As a negative control, the primary antibody was replaced by preimmune serum. The resulting staining was evaluated using a Zeiss confocal laser scanning microscope.

Statistical analyses

The results are presented as mean ± s.e.m. Overall analysis was carried out using the Kruskal–Wallis test, followed by the Mann–Whitney test for pairwise comparisons when overall significance was detected. The actual *N* value in the experiments that were performed to determine the effects of drugs on oviductal egg transport is the total number of rats used in each experimental group.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work was supported by FONDECYT (grant numbers 8980008, 1030315, 1080523, and 1040804), PROGRESAR (grant number PRE 004/2003), and Proyecto BASAL FBO-07.

Acknowledgements

We thank Andrea Gómez, PhD, from Millenium Nucleus of Stress and Addiction in Chile (NEDA) for figure design advice.

References

- Abudara V, Garcés G & Sáez JC 1999 Cells of the carotid body express connexin 43 which is up-regulated by cAMP. *Brain Research* **849** 25–33.
- Ardekani AM, Romanelli JC & Mayo KE 1998 Structure of the rat inhibin and activin betaA-subunit gene and regulation in an ovarian granulosa cell line. *Endocrinology* **139** 3271–3279.
- Bhatt P, Kadam K, Saxena A & Natraj U 2004 Fertilization, embryonic development and oviductal environment: role of estrogen induced oviductal glycoprotein. *Indian Journal of Experimental Biology* **42** 1043–1055.
- Bilezikjian LM, Corrigan AZ, Blount AL, Chen Y & Vale WW 2001 Regulation and actions of Smad7 in the modulation of activin, inhibin, and transforming growth factor-beta signaling in anterior pituitary cells. *Endocrinology* **142** 1065–1072.
- Bjornstrom L & Sjöberg M 2005 Mechanisms of estrogen receptor signaling: convergence of genomic and nongenomic actions on target genes. *Molecular Endocrinology* **19** 833–842.
- Blake CA, Brown LM, Duncan MW, Hunsucker SW & Helmke SM 2005 Estrogen regulation of the rat anterior pituitary gland proteome. *Experimental Biology and Medicine* **230** 800–807.
- Brann DW, Zhang QG, Wang RM, Mahesh VB & Vadlamudi RK 2008 PELP1 – a novel estrogen receptor-interacting protein. *Molecular and Cellular Endocrinology* **290** 2–7.
- Buhi WC 2002 Characterization and biological roles of oviduct-specific, oestrogen dependent glycoprotein. *Reproduction* **123** 355–362.
- Carlsson B, Hillensjö T, Nilsson A, Tornell J & Billig H 1993 Expression of insulin-like growth factor-I (IGF-I) in the rat fallopian tube: possible autocrine and paracrine action of fallopian tube-derived IGF-I on the fallopian tube and on the preimplantation embryo. *Endocrinology* **133** 2031–2039.
- Cheskis BJ, Greger JG, Nagpal S & Freedman LP 2007 Signaling by estrogens. *Journal of Cell Physiology* **213** 610–617.
- Cheskis BJ, Greger J, Cooch N, McNally C, McLaren S, Lam HS, Rutledge S, Mekonnen B, Hauze D, Nagpal S *et al.* 2008 MNAR plays an important role in ERα activation of Src/MAPK and PI3K/Akt signaling pathways. *Steroids* **73** 901–905.
- Clark BJ, Ranganathan V & Combs R 2001 Steroidogenic acute regulatory protein expression is dependent upon post-translational effects of cAMP-dependent protein kinase A. *Molecular and Cellular Endocrinology* **173** 183–192.
- Croxatto HB 1996 Gamete transport. In *Reproductive Endocrinology, Surgery and Technology*, edn 3, pp 385–402. Eds EY Adashi, JA Rock & Z Rosenwaks. Philadelphia: Lippincott-Raven Publishers.
- Croxatto HB 2002 Physiology of gamete and embryo transport through the fallopian tube. *Reproductive Biomedicine Online* **4** 160–169.
- Croxatto HB, Ortiz ME, Forcelledo ML, Fuentealba B, Noé G, Moore G, Moran F & Cárdenas H 1991 Hormonal control of ovum transport through the rat oviduct. *Archivos de Biología y Medicina Experimentales* **24** 403–410.
- Forcelledo ML, De la Cerda ML & Croxatto HB 1986 Effectiveness of different estrogen pulses in plasma for accelerating ovum transport and their relation to estradiol levels in the rat oviduct. *Endocrinology* **119** 1189–1194.
- Gómez-Sánchez EP, Ganjam V, Chen YJ, Liu Y, Zhou MY, Toroslu C, Romero DG, Hughson MD, de Rodríguez A & Gómez-Sánchez CE 2003 Regulation of 11 beta-hydroxysteroid dehydrogenase enzymes in the rat kidney by estradiol. *American Journal of Physiology. Endocrinology and Metabolism* **285** E272–E279.
- González-Arenas A, Villamar-Cruz O, Guerra-Araiza C & Camacho-Arroyo I 2003 Regulation of progesterone receptor isoforms expression by sex steroids in the rat lung. *Journal of Steroid Biochemistry and Molecular Biology* **85** 25–31.
- Hermoso M, Sáez JC & Villalón M 1997 Identification of gap junctions in the oviduct and regulation of connexins during development and by sexual hormones. *European Journal of Cell Biology* **74** 1–9.
- Ikeda Y, Nagai A, Ikeda MA & Hayashi S 2001 Neonatal estrogen exposure inhibits steroidogenesis in the developing rat ovary. *Developmental Dynamics* **221** 443–453.
- Irusta G, Parborell F, Peluffo M, Manna PR, Gonzalez-Calvar SI, Calandra R, Stocco DM & Tesone M 2003 Steroidogenic acute regulatory protein in ovarian follicles of gonadotropin-stimulated rats is regulated by a gonadotropin-releasing hormone agonist. *Biology of Reproduction* **68** 1577–1583.
- Jansen RP 1984 Endocrine response in the fallopian tube. *Endocrine Reviews* **5** 525–551.
- Livak KJ & Schmittgen TD 2001 Analysis of relative gene expression data using real-time quantitative PCR and the 2(T) (–Delta Delta C) method. *Methods* **25** 402–408.
- Mathieu CL, Mills SE, Burnett SH, Cloney DL, Bruns DE & Bruns ME 1989 The presence and estrogen control of immunoreactive calbindin-D9k in the fallopian tube of the rat. *Endocrinology* **125** 2745–2750.
- O'Lone R, Frith MC, Karlsson EK & Hansen U 2004 Genomic targets of nuclear estrogen receptors. *Molecular Endocrinology* **18** 1859–1875.
- Orihuela PA & Croxatto HB 2001 Acceleration of oviductal transport of oocytes induced by estradiol in cycling rats is mediated by nongenomic stimulation of protein phosphorylation in the oviduct. *Biology of Reproduction* **65** 1238–1245.
- Orihuela PA, Ríos M & Croxatto HB 2001 Disparate effects of estradiol on egg transport and oviductal protein synthesis in mated and unmated rats. *Biology of Reproduction* **65** 1232–1237.

- Orihuela PA, Parada-Bustamante A, Cortés PP, Gatica C & Croxatto HB** 2003 Estrogen receptor, cyclic adenosine monophosphate, and protein kinase a are involved in the nongenomic pathway by which estradiol accelerates oviductal oocyte transport in cyclic rats. *Biology of Reproduction* **68** 1225–1231.
- Orihuela PA, Zuñiga LM, Ríos M, Lladós C, Sierralta W & Croxatto HB** 2004 Estrogen receptors in the rat oviduct. *Molecular Biology of the Cell* **15** 265A.
- Orihuela PA, Parada-Bustamante A, Zuñiga LM & Croxatto HB** 2006 Inositol triphosphate participates in an oestradiol nongenomic signalling pathway involved in accelerated oviductal transport in cycling rats. *Journal of Endocrinology* **188** 579–588.
- Ortiz ME, Villalón M & Croxatto HB** 1979 Ovum transport and fertility following post-ovulatory treatment with estradiol in rats. *Biology of Reproduction* **21** 1163–1167.
- Parada-Bustamante A, Orihuela PA & Croxatto HB** 2003 Effect of intrauterine insemination with spermatozoa or foreign protein on the mechanism of action of oestradiol in the rat oviduct. *Reproduction* **125** 677–682.
- Parada-Bustamante A, Orihuela PA, Ríos M, Navarrete-Gómez PA, Cuevas CA, Velásquez LA, Villalón MJ & Croxatto HB** 2007 Catechol-O-methyltransferase and methoxyestradiols participate in the intraoviductal nongenomic pathway through which estradiol accelerates egg transport in cycling rats. *Biology of Reproduction* **77** 934–941.
- Peñarroja-Matutano C, Parada-Bustamante A, Orihuela PA, Ríos M, Velásquez LA & Croxatto HB** 2007 Genital sensory stimulation shifts estradiol intraoviductal signaling from nongenomic to genomic pathways, independently from prolactin surges. *Biological Research* **40** 213–222.
- Pérez Martínez S, Hermoso M, Farina M, Ribeiro ML, Rapanelli M, Espinosa M, Villalón M & Franchi A** 2006 17 β -Estradiol upregulates COX-2 in the rat oviduct. *Prostaglandins & Other Lipid Mediators* **80** 155–164.
- Priyadarsana M, Wijayagunawardane B & Miyamoto A** 2004 Endothelin-1 system in the bovine oviduct: a regulator of local contraction and gamete transport. *Journal of Cardiovascular Pharmacology* **44** S248–S251.
- Ríos M, Orihuela PA & Croxatto HB** 1997 Intraoviductal administration of ribonucleic acid from estrogen-treated rats mimics the effect of estrogen on ovum transport. *Biology of Reproduction* **56** 279–283.
- Ríos M, Hermoso M, Sánchez TM, Croxatto HB & Villalón MJ** 2007 Effect of oestradiol and progesterone on the instant and directional velocity of microsphere movements in the rat oviduct: gap junctions mediate the kinetic effect of oestradiol. *Reproduction, Fertility, and Development* **19** 634–640.
- Scott RE, Wu-Peng XS, Kaplitt MG & Pfaff DW** 2003 Gene transfer and *in vivo* promoter analysis of the rat progesterone receptor using a herpes simplex virus viral vector. *Brain Research. Molecular Brain Research* **114** 91–100.
- Smith MS, Freeman ME & Neill JD** 1975 The control of progesterone secretion during the estrous cycle and early pseudopregnancy in the rat: prolactin, gonadotropin and steroid levels associated with rescue of the corpus luteum of pseudopregnancy. *Endocrinology* **96** 219–226.
- Stocco DM & Clark BJ** 1996 Regulation of the acute production of steroids in steroidogenic cells. *Endocrine Reviews* **17** 221–244.
- Stocco DM, Wang X, Jo Y & Manna PR** 2005 Multiple signaling pathways regulating steroidogenesis and steroidogenic acute regulatory protein expression: more complicated than we thought. *Molecular Endocrinology* **19** 2647–2659.
- Turner CD** 1961 Endocrinology of the ovary. In *General Endocrinology*, edn 3, pp 365–409. Ed. CD Turner. Philadelphia: WB Saunders Company.
- Villalón M & Verdugo P** 1982 Hormonal regulation of ciliary function in the oviduct: the effect of beta-adrenergic agonists. *Progress in Clinical and Biological Research* **80** 59–65.
- Ye F, Florian M, Magder SA & Hussain SN** 2004 Regulation of angiotensin and Tie-2 receptor expression in non-reproductive tissues by estrogen. *Steroids* **67** 305–310.

Received 29 May 2009

First decision 24 August 2009

Revised manuscript received 10 November 2009

Accepted 23 December 2009